



I hereby certify that this correspondence is being transmitted via first class mail to the Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on:

26 Dec. 2003

Date of Deposit

Kathryn A. Kallay

Signature of Person Mailing Paper

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re David Alexander Learmonth et al.

Serial No. 10/681578

Confirmation No.

Filed: 8 October 2003

For: PERIPHERALLY-SELECTIVE INHIBITORS
OF DOPAMINE- β -HYDROXYLASE AND
METHOD OF THEIR PREPARATION

Group Art Unit:

Examiner:

Attorney Docket: AAT-15033

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

**SUBMISSION OF PRIORITY DOCUMENTS AND
REAFFIRMATION OF PRIORITY CLAIM**

Sir:

Applicants hereby reaffirm their priority claim in the above noted application to the following documents for which certified copies are included herewith:

	<u>Application No.</u>	<u>Application Date</u>
1. United Kingdom	0223719.6	October 11, 2002
2. United Kingdom	0224306.1	October 18, 2002

Respectfully submitted,

RANKIN, HILL, PORTER & CLARK LLP

Kenneth A. Clark, Reg. No. 32,119

925 Euclid Avenue, Suite 700
Cleveland, Ohio 44115-1405
(216) 566-9700





INVESTOR IN PEOPLE


The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.




P. Mahoney

Signed

Dated 3 November 2003





Request for grant of a patent

(See the notes on the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form)

The Patent Office

Cardiff Road
Newport
South Wales
NP9 1RH

1. Your reference

PAC/20930 GB

2. Patent application number

(The Patent Office will fill in this part)

0223719.6

3. Full name, address and postcode of the or of each applicant (underline all surnames)

PORTELA & C.A., S.A.

À AV. DA SIDERURGIA NACIONAL
APARTADO 19

4745-457 S. MAMEDE DO CORONADO
PORTUGAL

Patents ADP number (if you know it)

If the applicant is a corporate body, give the country/state of its incorporation

PORTUGAL

7572423002

4. Title of the invention

PERIPHERALLY-SELECTIVE INHIBITORS
OF DOPAMINE-BETA-HYDROXYLASE AND
METHOD OF THEIR PREPARATION

5. Name of your agent (if you have one)

A A THORNTON & CO

"Address for service" in the United Kingdom
to which all correspondence should be sent
(including the postcode)

235 HIGH HOLBORN
LONDON WC1V 7LE

Patents ADP number (if you know it)

0000075001

6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number

Country

Priority application number
(if you know it)

Date of filing
(day / month / year)

7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application

Number of earlier application

Date of filing
(day / month / year)

8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer 'Yes' if:

YES

- a) any applicant named in part 3 is not an inventor, or
 - b) there is an inventor who is not named as an applicant, or
 - c) any named applicant is a corporate body.
- See note (d))

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description

23 ✓

Claim(s)

6 ✓

Abstract

1 ✓

Drawing(s)

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

YES ✓

Request for preliminary examination and search (Patents Form 9/77)

YES ✓

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application.

Signature



Date

A. A. Thornton & Co.

11 OCTOBER 2002

12. Name and daytime telephone number of person to contact in the United Kingdom

Philip A. CURTIS - 0207 440 6860

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

- If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.
- Write your answers in capital letters using black ink or you may type them.
- If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.
- If you have answered 'Yes' Patents Form 7/77 will need to be filed.
- Once you have filled in the form you must remember to sign and date it.
- For details of the fee and ways to pay please contact the Patent Office.

**PERIPHERALLY-SELECTIVE INHIBITORS OF DOPAMINE- β -HYDROXYLASE AND
METHOD OF THEIR PREPARATION**

5 This invention relates to peripherally-selective inhibitors of dopamine- β -hydroxylase and method of their preparation.

In recent years, interest in the development of inhibitors of dopamine- β -hydroxylase (D β H) has centred on the hypothesis that inhibition of this enzyme may provide
10 significant clinical improvements in patients suffering from cardiovascular disorders such as hypertension or chronic heart failure. The rationale for the use of D β H inhibitors is based on their capacity to inhibit the biosynthesis of noradrenaline, which is achieved via enzymatic hydroxylation of dopamine. Activation of neurohumoral systems, chiefly the sympathetic nervous system, is the principal clinical manifestation of congestive
15 heart failure (Parmley, W.W., Clinical Cardiology, 18: 440-445, 1995). Congestive heart failure patients have elevated concentrations of plasma noradrenaline (Levine, T.B. et al., Am. J. Cardiol., 49:1659-1666, 1982), increased central sympathetic outflow (Leimbach, W.N. et al., Circulation, 73: 913-919, 1986) and augmented cardiorenal noradrenaline spillover (Hasking, G.J. et al., Circulation, 73:615-621, 1966). Prolonged
20 and excessive exposure of myocardium to noradrenaline may lead to down-regulation of cardiac β_1 -adrenoceptors, remodelling of the left ventricle, arrhythmias and necrosis, all of which can diminish the functional integrity of the heart. Congestive heart failure patients who have high plasma concentrations of noradrenaline also have the most unfavourable long-term prognosis (Cohn, J.N. et al., N. Engl. J. Med., 311:819-823,
25 1984). Of greater significance is the observation that plasma noradrenaline concentrations are already elevated in asymptomatic patients with no overt heart failure and can predict ensuing mortality and morbidity (Benedict, C.R. et al., Circulation, 94:690-697, 1996). This implies that the activated sympathetic drive is not merely a clinical marker of congestive heart failure, but may contribute to progressive worsening
30 of the disease.

Inhibition of sympathetic nerve function with adrenoceptor antagonists appeared a promising approach, however a significant proportion of patients do not tolerate the

immediate haemodynamic deterioration that accompanies β -blocker treatment (Pfeffer, M.A. et al., N. Engl. J. Med., 334:1396-7, 1996). An alternative strategy for directly modulating sympathetic nerve function is to reduce the biosynthesis of noradrenaline via inhibition of D β H, the enzyme responsible for conversion of dopamine to noradrenaline in sympathetic nerves. This approach has several merits including gradual modulation as opposed to abrupt inhibition of the sympathetic system, and causing increased release of dopamine, which can improve renal function such as renal vasodilation, diuresis and natriuresis. Therefore inhibitors of D β H may provide significant advantages over conventional β -blockers.


10

Several inhibitors of D β H have been thus far reported in the literature. Early first and second generation examples such as disulfiram (Goldstein, M. et al., Life Sci., 3:763, 1964) and diethyldithiocarbamate (Lippmann, W. et al., Biochem. Pharmacol., 18: 2507, 1969) or fusaric acid (Hidaka, H. Nature, 231, 1971) and aromatic or alkyl thioureas (Johnson, G.A. et al, J. Pharmacol. Exp. Ther., 171: 80, 1970) were found to be of low potency, exhibited poor selectivity for D β H and caused toxic side effects. The third generation of D β H inhibitors however, were found to have much greater potency, such as for example, nepicastat (RS-25560-197, IC₅₀ 9nM) (Stanley, W.C., et al., Br. J Pharmacol., 121: 1803-1809, 1997), which was developed to early clinical trials. Although devoid of some of the problems associated with first and second generation D β H inhibitors, a very important discovery was that nepicastat was found to cross the blood brain barrier (BBB), thereby able to cause central as well as peripheral effects, a situation which could lead to undesired and potentially serious CNS side-effects of the drug. Therefore there yet remains an unfulfilled clinical requirement for a potent, non-toxic and peripherally selective inhibitor of D β H, which could be used for treatment of certain cardiovascular disorders. A D β H inhibitor with similar or even greater potency than nepicastat, but devoid of CNS effects (inability to cross the BBB) would provide a significant improvement over all D β H inhibitor compounds thusfar described in the prior art.

30

We have surprisingly found that incorporation of certain heteroatoms to the carbocyclic ring and/or elongation of the amino alkyl side-chain of the nepicastat core-structure

gives rise to a series of compounds possessing significant and pronounced effects of potential usefulness for D β H inhibition. Many of these compounds are endowed with greater potency and significantly reduced brain access, giving rise to potent and peripherally selective D β H inhibitors. Thus, the invention relates to compounds of
5 general formula I;




I

where R₁, R₂ and R₃ are the same or different and signify hydrogens, halogens, alkyl, alkylaryl, alkyloxy, hydroxy, nitro, amino, alkylcarbonylamino, alkylamino or
10 dialkylamino group; R₄ signifies hydrogen, alkyl or alkylaryl group; X signifies CH₂, oxygen atom or sulphur atom; n is 1, 2 or 3, with the proviso that when n is 1, X is not CH₂; and the individual (R)- and (S)-enantiomers or mixtures of enantiomers; and the pharmaceutically acceptable salts thereof.

15 Unless stated otherwise, in this specification the term alkyl (whether used on its own or used in combination with other moieties) means hydrocarbon chains, straight or branched, containing from one to six carbon atoms, optionally substituted by aryl, alkoxy, halogen, alkoxycarbonyl or hydroxycarbonyl groups; the term aryl (whether used on its own or used in combination with other moieties) means a phenyl or
20 naphthyl group, optionally substituted by alkyloxy, halogen or nitro group; and the term halogen means fluorine, chlorine, bromine or iodine.


Another aspect of the present invention is a process for the preparation of compounds of formula I. Some compounds according to formula II where X signifies methylene
25 (CH₂), oxygen or sulphur are known (Martinez, G.R. et al., US Patent 5,538,988, Jul. 23, 1996; Eriksson, M., PCT Int. Appl. WO 9959988A1, 25 Nov 1999; Napoletano, M., PCT Int. Appl. WO 9608489A1, 21 March 1996; Sarda, N. et al., Tetrahedron Lett., 17:271-272, 1976; Neirabeyeh, M.AI et al., Eur. J. Med. Chem., 26:497-504, 1991) in

the literature and others can be prepared by those skilled in the art. Compounds according to formula II are chiral, and formula II is therefore to be taken to represent both optically pure individual (R)- and (S)-enantiomers or mixtures of enantiomers;



II

- 5 Compounds of formula I are prepared by reacting a compound of formula II where X is CH₂, oxygen or sulphur; R₁, R₂ and R₃ are the same or different and signify hydrogens, halogens, alkyl, alkylaryl, alkyloxy, hydroxy, nitro, alkylcarbonylamino, alkylamino or dialkylamino group with a compound of formula III:



III

10

- where n signifies 1, 2 or 3; when n is 1 or 2, R₄ signifies hydrogen, alkyl or alkylaryl group; R₅ signifies a hydroxyl protecting group and R₆ signifies an amino protecting group; when n signifies 3, R₅ is defined as above but R₄ and R₆ taken together represent a phthalimido group; and with a water soluble thiocyanate salt in an inert
 15 organic solvent and in the presence of an organic acid, wherein the water soluble thiocyanate salt is an alkali metal thiocyanate salt or a tetraalkylammonium thiocyanate salt.

- Suitable alkali metal thiocyanate salts include sodium, lithium and cesium thiocyanates,
 20 but potassium thiocyanate is preferred.

- The compound of formula III where n is 1 is known (Wolf, E. et al., Can. J. Chem., 75:942-948, 1997) and compounds of formula III where n is 2 or 3 are new compounds that can be prepared by those skilled in the art (see examples). The preferred hydroxyl
 25 protecting groups (R₅) include organosilyl compounds such as chosen from trialkylsilyl, triphenylsilyl, phenyldialkylsilyl or alkyldiphenylsilyl group. The tert-butyldimethylsilyl (TBDMS) group is especially preferred. The preferred amino protecting groups (R₆)

include carbamates such alkyl carbamates, in particular the t-butyl carbamate (Boc) group, and alkylaryl carbamates. The reaction may be run with a small excess of the compound of formula III and potassium thiocyanate (preferably 1.1-1.3 equivalents).

- 5 The invention also provides compounds of formula II, where at least one of R_1 , R_2 and R_3 is fluorine.


The reaction can be run in a substantially inert solvent (preferably ethyl acetate) and at different temperatures (preferably at the solvent reflux temperature). Preferred organic
10 acids include acetic acid. When compounds of formula III where n signifies 1 are used, the intermediate of formula IV is then treated with a mineral acid in a suitable solvent to remove the Boc amino protecting group and provide the compounds of formula I (scheme 1). Preferred mineral acids include hydrochloric acid and preferred solvents include ethyl acetate.

15


When compounds of formula III where n signifies 2 are used and R_4 signifies hydrogen, the mixture of intermediate products of formula V and VI is reacted with hydrochloric acid in ethyl acetate to afford the corresponding single compounds of formula I (scheme 2); where R_4 signifies alkyl (including alkyl substituted by aryl), the single
20 intermediate product of formula V is reacted with hydrochloric acid in ethyl acetate to afford the compounds of formula I.

When compounds of formula III where n is 3 are used, the intermediate of formula VII is then treated with sodium borohydride in a suitable solvent system followed by acetic
25 acid to remove the pthalimido amino protecting group as described in the literature (Osby et al., Tetrahedron Lett., 1984, 25(20), 2093-2096) to give the compounds of formula I (scheme 3). The compounds of formula I are obtained with good purity, but if preferred can be recrystallised from a suitable solvent.

5


Scheme 1

Scheme 2



Scheme 3

5



For the preparation of pharmaceutical compositions of compounds of formula I, inert pharmaceutically acceptable carriers are admixed with the active compounds. The pharmaceutically acceptable carriers may be either solid or liquid. Solid form preparations include powders, tablets, dispersible granules and capsules. A solid carrier can be one or more substances which may also act as diluents, flavouring agents, solubilizers, lubricants, suspending agents, binders or tablet disintegrating agents; it may also be an encapsulating material.

Preferably the pharmaceutical preparation is in unit dosage form, e.g. packaged preparation, the package containing discrete quantities of preparation such as packeted tablets, capsules and powders in vials or ampoules.

The dosages may be varied depending on the requirement of the patient, the severity of the disease and the particular compound being employed. For convenience, the total daily dosage may be divided and administered in portions throughout the day. It is expected that once or twice per day administration will be most suitable. Determination of the proper dosage for a particular situation is within the skill of those in the medical art.

20 Materials and Methods

In vitro studies

D β H activity was evaluated by the ability to β -hydroxylate dopamine to noradrenaline as previously described (Kojima, K., Parvez, S. and Nagatsu T. 1993. Analysis of enzymes in catecholamine biosynthesis. In Methods in Neurotransmitter and Neuropeptide Research, pp. 349-380: Elsevier Science Publishers). SK-N-SH cells (ATCC HTB-11), a human neuroblastoma derived cell line, were used as a source of human D β H. SK-N-SH cells cultured in 24 well plates were preincubated for 20 min in a reaction medium containing 200 mM sodium acetate, 30 mM *N*-ethylmaleimide, 5 μ M copper sulphate, 0.5 mg/ml catalase aqueous solution, 1 mM pargyline, 10 mM sodium fumarate and 20 mM ascorbic acid. Thereafter, cells were incubated for further 45 min in the reaction medium with added increasing concentrations of dopamine (0.5 to 100 mM). During

preincubation and incubation, the cells were continuously shaken and maintained at 37°C. The reaction was terminated by the addition of 0.2 M perchloric acid. The acidified samples were stored at 4°C before injection into the high pressure liquid chromatograph for the assay of noradrenaline. In experiments conducted with the aim of studying the effects of new D β H inhibitors on enzyme activity, test compounds (0.3 to 10,000 nM) of interest were added to the preincubation and incubation solutions; the incubation was performed in the presence of a concentration (50 mM) of dopamine 2.5 times the corresponding K_m value as determined in saturation experiments.

10 *In vivo* studies

Male NMRI mice or Wistar rats were obtained from Harlan-Interfauna (Spain) and were kept 10 and 5 per cage respectively, under controlled environmental conditions (12 h light/dark cycle and room temperature 22 ± 1 °C). Food and tap water were allowed *ad libitum* and experimentation was performed during daylight hours.

At time = 0 h, animals were administered with either test compounds at a given dose or vehicle (water) delivered orally *via* gavage. At 2, 6, 9, 12, 18 and 24 h post dose, the animals were sacrificed by decapitation and heart (left atrium and left ventricle) and brain (frontal and parietal cortex) were isolated, weighed and stored in a volume of 0.2 M perchloric acid for 12 h at 4 °C in the dark. Post incubation, the resulting supernatants were collected by centrifuge filtration of incubates (0.2 μ M / 10 min / ~5000 rpm, 4 °C). Supernatants were stored frozen at -80 °C until analysis. Quantification of dopamine and noradrenaline in supernatants was performed by high pressure liquid chromatography with electrochemical detection.

R sults

In vitro studies

5

Incubation of SK-N-SH cells in the presence of increasing concentrations of dopamine resulted in a concentration-dependent formation of noradrenaline, yielding K_m (in μM) and V_{max} (in $nmol\ mg\ protein^{-1}\ h^{-1}$) values of 20.6 ± 1.6 and 153.8 ± 4.4 , respectively. From these kinetic parameters, a concentration of dopamine approaching saturation 10 (50 mM) was chosen for use in inhibition studies. As listed in Table 1 compounds **2,3,4,5,6,7,8,10,12,16,19,24,26,28** and **29** were found to markedly inhibit D β H activity. Compounds **2**, **3**, **4** and nepicastat **1** (the reference compound) produced a concentration-dependent decrease in the β -hydroxylation of dopamine with IC_{50} values in the low nM range against human D β H activity (see Table 2). Compound **4** was 15 chosen for further *in vivo* studies, being the compound most closely related to nepicastat **1** in order to provide conclusive evidence that the structural modifications made to the molecule as part of the present invention are responsible for the surprisingly markedly improved biological properties observed.

Table 1. Effect of selected compounds (5 μ M) on D β H activity in SK-N-SH cells. Values are quoted as % of control.

No.	Mean $\pm$ SEM	No.	Mean $\pm$ SEM
1	0.0 $\pm$ 0.3	24	0.0 $\pm$ 1.9
2	1.6 $\pm$ 0.3	25	66.0 $\pm$ 4.5
3	4.1 $\pm$ 0.6	26	4.5 $\pm$ 1.9
4	3.3 $\pm$ 0.3	27	15.5 $\pm$ 5.8
5	8.1 $\pm$ 0.3	28	2.6 $\pm$ 1.6
6	6.9 $\pm$ 0.6	29	2.2 $\pm$ 2.5
7	8.0 $\pm$ 0.1	30	99.4 $\pm$ 2.8
8	9.4 $\pm$ 0.7	31	27.3 $\pm$ 0.4
9	50.2 $\pm$ 1.9		
10	8.2 $\pm$ 0.7		
11	36.7 $\pm$ 4.4		
12	3.0 $\pm$ 0.5		
13	94.0 $\pm$ 3.1		
14	77.9 $\pm$ 2.2		
15	86.1 $\pm$ 2.7		
16	0.0 $\pm$ 0.6		
17	53.2 $\pm$ 3.9		
18	94.8 $\pm$ 1.2		
19	6.9 $\pm$ 0.5		
20	16.8 $\pm$ 4.8		
21	124.8 $\pm$ 6.5		
22	17.8 $\pm$ 2.1		
23	54.5 $\pm$ 9.9		

5

Table 2. IC₅₀ values (in nM) for inhibition of D β H in SK-N-SH cells.

Compound	IC ₅₀ (in nM)
2	60 (14, 250)
3	91 (56, 147)
4	105 (69, 161)
Nepicastat 1	36 (28, 46)

10

In vivo studies

Mouse

5 The time course experiments for compound **4** and nepicastat (**1**) in the heart at 100 mg/kg suggests that both compounds are long acting. Time of maximum effect ($T_{\max}$) for noradrenaline tissue reduction by both **4** and **1** appears to be at 9 h post-dose (Figure 1). Thereafter, noradrenaline tissue levels recover, reaching 50% recovery of initial tissue levels at 24 h.

10

At $T_{\max}$ (9 h after administration), both **4** and **1** reduced noradrenaline levels in a dose-dependent manner in left ventricle. For both **4** and **1**, the maximal inhibitory effect was attained at a dose of 100 mg/kg. In contrast to that found in the heart, **4** failed to affect noradrenaline tissue levels in the brain parietal cortex, whereas **1** produced a dose-
15 dependent decrease in noradrenaline levels in this area of the brain (Figure 2).

Rat

As shown in the mouse, the effects of both **4** and **1** upon noradrenaline were
20 dependent on the dose administered and reached its maximum at 9 h (data not shown). However, as depicted in Figure 3, the inhibitory effects of **4** (100 mg/kg) upon noradrenaline levels in both the left atrium and the left ventricle were more pronounced than those elicited by **1** (100 mg/kg). Again, as observed in the mouse, **4** failed to affect noradrenaline tissue levels in the brain parietal cortex and the brain frontal cortex,
25 whereas **1** produced a marked decrease in noradrenaline levels in these brain areas.

It is concluded that **4**, in stark contrast to nepicastat **1**, exerts its inhibitory effects upon DBH exclusively in the periphery, being devoid of inhibitory effects in the brain.

30 Reference is now made to the accompanying drawings, in which:

Figure 1 is a graph showing the time-dependent decrease of noradrenaline levels in the left ventricle of mice treated orally with 100 mg/kg of **4** or nepicastat **1**. Symbols are means of 5 determinations per group; vertical lines indicate S.E.M.

Figure 2 is two graphs showing noradrenaline levels in the mouse left ventricle and brain parietal cortex 9 h after oral administration of **4** or nepicastat **1**. Symbols are means of 5 determinations per group; vertical lines indicate S.E.M.

5

Figure 3 is four graphs showing noradrenaline levels in the rat heart (left atrium and left ventricle) and brain (frontal and parietal cortex) 9 h after the oral administration of **4** or nepicastat **1**. Columns are means of 5 determinations per group; vertical lines indicated S.E.M.

10

Conclusion

Some compounds of general formula I are very potent dopamine- β -hydroxylase inhibitors and have potentially valuable pharmaceutical properties in the treatment of some cardiovascular disorders, where a reduction in the enzymatic hydroxylation of dopamine to noradrenaline may be of therapeutic benefit, such as hypertension and chronic heart failure. The possibility to use a long-acting D β H inhibitor with limited access to the brain (CNS), such as compound **4** opens new perspectives in the treatment of hypertension and chronic heart failure by improving potency and selectivity of D β H inhibition in the periphery.

20

The invention disclosed herein is exemplified by the following examples of preparation, which should not be construed to limit the scope of the disclosure. Alternative pathways and analogous structures may be apparent to those skilled in the art.

25 Examples

Example 1

(*R*)-5-aminomethyl-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione

30 hydrochloride compound **3**, table 1)

A stirred mixture of (*R*)-6,8-difluorochroman-3-ylamine hydrochloride (0.22 g, 1.0 mmol), [3-(tert-butyl(dimethyl)silyloxy)-2-oxopropyl]carbamic acid tert-butyl ester (0.33 g, 1.1 mmol), potassium thiocyanate (0.11 g, 1.1 mmol) and acetic acid (0.3 mL,

- 5.0 mmol) in ethyl acetate (3 mL) was refluxed for 2 hours, cooled to room temperature, then washed by sodium bicarbonate solution, dried over anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by the column chromatography over silica gel using ethyl acetate – petroleum ether mixture as eluent.
- 5 The resulting oil (0.23 g) was dissolved in ethyl acetate (2 ml), whereupon 2M HCl solution in ethyl acetate was added (2 mL, 4 mmol) and the mixture was stirred for 2 hours at room temperature. The precipitate was removed by filtration and washed with ethyl acetate to give crystals of m.p. 192 °C (decomp.).

10 Examples 2-3

By the application of the above described technique and related procedures known to those skilled in the art and using the appropriate chroman-3-ylamines hydrochlorides, the following compounds were prepared:

- 15 (*R*)-5-aminomethyl-1-chroman-3-yl-1,3-dihydroimidazole-2-thione hydrochloride (compound **24**, table 1)
- (*R*)-5-aminomethyl-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **22**, table 1)

Example 4

(*R,S*)-5-aminomethyl-1-(6-hydroxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride

- 5 A stirred mixture of 6-hydroxythiochroman-3-ylamine hydrochloride (0.22 g, 1.0 mmol), [3-(tert-butyldimethylsilyloxy)-2-oxopropyl]carbamic acid tert-butyl ester (0.33 g, 1.1 mmol), potassium thiocyanate (0.11 g, 1.1 mmol) and acetic acid (0.3 mL, 5.0 mmol) in ethyl acetate (3 mL) was refluxed for 2 hours, then cooled to room temperature, and washed by sodium bicarbonate solution, dried over anhydrous
- 10 magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.25 g) was dissolved in ethyl acetate (2 mL), whereupon 2M HCl solution in ethyl acetate was added (2 mL, 4 mmol) and the mixture was stirred for 2 hours at room temperature. The precipitate was removed by filtration and washed with ethyl
- 15 acetate to give crystals, which decomposed without melting.

Example 5

(3,4-Dihydroxybutyl)carbamic acid tert-butyl ester

- 20 To a stirred solution of 4-amino-1,2-propanediol (2.10 g, 20 mmol) in ethanol (50 mL) at room temperature was added di-tert-butyldicarbonate (4.80 g, 22 mmol) in one portion. The resulting mixture was stirred at room temperature for two hours, then evaporated in vacuo and purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent to afford colourless oil.

25

Examples 6-7

- By the application of the above described technique and related procedures known to those skilled in the art and using the appropriate N-substituted 4-amino-1,2-
- 30 propanediols, the following compounds were prepared:

(3,4-Dihydroxybutyl)methylcarbamic acid tert-butyl ester

(3,4-Dihydroxybutyl)benzylcarbamic acid tert-butyl ester

Example 8

[4-(tert-butyldimethylsilyloxy)-3-hydroxybutyl]carbamic acid tert-butyl ester

To a stirred solution of (3,4-dihydroxybutyl)carbamic acid tert-butyl ester (2.60 g, 12.7 mmol), triethylamine (2.03 mL, 14.50 mmol) and 4-(dimethylamino)pyridine (0.05 g, 0.4 mmol) in anhydrous dichloromethane (40 mL) at room temperature was added tert-butyldimethylchlorosilane (2.0 g, 13.17 mmol) in one portion. The resulting mixture was stirred at room temperature for 18 hours, washed with water, brine and dried over anhydrous magnesium sulfate. Filtration and concentration in vacuo gave an oil which was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent to afford a colourless oil.

Example 9-10

By the application of the above described technique and related procedures known to those skilled in the art and using compounds from examples 6 and 7, the following compounds were prepared:

[4-(tert-butyldimethylsilyloxy)-3-hydroxybutyl]methylcarbamic acid tert-butyl ester

[4-(tert-butyldimethylsilyloxy)-3-hydroxybutyl]benzylcarbamic acid tert-butyl ester

Example 11

[4-(tert-butyldimethylsilyloxy)-3-oxobutyl]carbamic acid tert-butyl ester

To a solution of Dess-Martin periodinane (5.0 g, 11.8 mmol) in anhydrous dichloromethane (35 mL) at room temperature was added a solution of [4-(tert-butyldimethylsilyloxy)-3-hydroxybutyl]carbamic acid tert-butyl ester (3.77 g, 11.8 mmol) in anhydrous dichloromethane. The resulting mixture was stirred at room temperature for one hour, evaporated in vacuo to one third of the initial volume and applied to a column packed with silica. Elution with ethyl acetate – petroleum ether solvent mixture afforded a colourless oil.

Example 12-13

By the application of the above described technique and related procedures known to those skilled in the art and using compounds from examples 9 and 10, the following compounds were prepared:

[4-(tert-butyldimethylsilanyloxy)-3-oxobutyl]methylcarbamic acid tert-butyl ester
[4-(tert-butyldimethylsilanyloxy)-3-oxobutyl]benzylcarbamic acid tert-butyl ester.

Example 14

10

(S)-5-(2-aminoethyl)-1-(5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride, compound 2, table 1)

A stirred mixture of (S)-5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl amine hydrochloride (0.17 g, 0.79 mmol), [4-(tert-butyldimethylsilanyloxy)-3-oxobutyl]carbamic acid tert-butyl ester (0.28 g, 0.87 mmol), potassium thiocyanate (0.085 g, 0.85 mmol), water (0.014 mL, 0.80 mmol) and acetic acid (0.2 mL, 3.3 mmol) in ethyl acetate (2 mL) was refluxed for 7 hours, cooled to the room temperature, washed by sodium bicarbonate solution and dried over anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.24 g) was dissolved in ethyl acetate (2 mL), 2M HCl solution in ethyl acetate was added (2 mL, 4 mmol) and the mixture was stirred for 2 hours at room temperature. The precipitate was removed by filtration and washed with ethyl acetate to give crystals, which decomposed without melting.

25

Example 15

By the application of the above described technique and related procedures known to those skilled in the art and using the appropriate 1,2,3,4-tetrahydronaphthalen-2-ylamines hydrochlorides, the following compounds were prepared:

(S)-5-(2-aminoethyl)-1-(1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 20, table 1)

Example 16

(*R*)-5-(2-aminoethyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 4, table 1)

5 A stirred mixture of (*R*)-6,8-difluorochroman-3-ylamine hydrochloride (1.68 g, 7.58 mmol), [4-(tert-butyldimethylsilyloxy)-3-oxobutyl]carbamic acid tert-butyl ester (3.13 g, 9.85 mmol), potassium thiocyanate (0.96 g, 9.85 mmol), water (0.18 mL, 10 mmol) and acetic acid (3.0 mL, 50 mmol) in ethyl acetate (30 mL) was refluxed for 7 hours, cooled to room temperature, washed by sodium bicarbonate solution, dried over
 10 anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (2.15 g) was dissolved in ethyl acetate (20 mL), 2M HCl solution in ethyl acetate was added (20 mL, 40 mmol) and the mixture was stirred for 2 hours at room temperature. The precipitate was removed by filtration and washed with ethyl
 15 acetate to give crystals, which decomposed without melting.

Example 17-37

By the application of the above described technique and related procedures
 20 known to those skilled in the art and using the appropriate chroman-3-ylamine hydrochlorides and [4-(tert-butyldimethylsilyloxy)-3-oxobutyl]carbamic acid tert-butyl esters, the following compounds were prepared:

(*R*)-5-(2-aminoethyl)-1-chroman-3-yl-1,3-dihydroimidazole-2-thione hydrochloride (compound 12, table 1)

25 (*R*)-5-(2-aminoethyl)-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 16, table 1)

(*R*)-5-(2-aminoethyl)-1-(8-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 21, table 1)

(*R*)-5-(2-aminoethyl)-1-(6-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione
 30 hydrochloride (compound 23, table 1)

(*R*)-5-(2-aminoethyl)-1-(8-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 19, table 1)

(*R*)-5-(2-aminoethyl)-1-(6-fluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 7, table 1)

- (*R*)-5-(2-aminoethyl)-1-(8-fluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **6**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6,7-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **8**, table 1)
- 5 (*S*)-5-(2-aminoethyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **9**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6,7,8-trifluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **10**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6-chloro-8-methoxychroman-3-yl)-1,3-dihydroimidazole-2-
- 10 thione hydrochloride (compound **11**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6-methoxy-8-chlorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **13**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6-nitrochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **18**, table 1)
- 15 (*R*)-5-(2-aminoethyl)-1-(8-nitrochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **17**, table 1)
- (*R*)-5-(2-aminoethyl)-1-[6-(acetylamino)chroman-3-yl]-1,3-dihydroimidazole-2-thione hydrochloride (compound **14**, table 1)
- (*R*)-5-(2-aminoethyl)-1-(6-hydroxy-7-benzylchroman-3-yl)-1,3-dihydroimidazole-2-thione
- 20 hydrochloride (compound **15**, table 1)
- (*R*)-5-(2-Benzylaminoethyl)-1-(6-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **25**, table 1)
- (*R*)-5-(2-Benzylaminoethyl)-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **26**, table 1)
- 25 (*R*)-1-(6-Hydroxychroman-3-yl)-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **27**, table 1)
- (*R*)-1-(6,8-Difluorochroman-3-yl)-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **28**, table 1)
- (*R*)-1-Chroman-3-yl-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione
- 30 hydrochloride (compound **29**, table 1)

Example 38

(*R,S*)-5-(2-aminoethyl)-1-(6-methoxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **30**, table 1)

A stirred mixture of 6-methoxythiochroman-3-ylamine hydrochloride (0.12 g, 0.50 mmol), [3-(tert-butyldimethylsilyloxy)-2-oxopropyl]carbamic acid tert-butyl ester (0.17 g, 0.55 mmol), potassium thiocyanate (0.055 g, 0.55 mmol), water (0.009 g, 0.50 mmol) and acetic acid (0.2 mL, 3.3 mmol) in ethyl acetate (2 mL) was refluxed for 7 hours, cooled to room temperature, washed by sodium bicarbonate solution, dried over anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.12 g) was dissolved in ethyl acetate (1 mL), 2M HCl solution in ethyl acetate was added (1 mL, 2 mmol) and the mixture was stirred for 2 hours at room temperature. The precipitate was removed by filtration and washed with ethyl acetate to give crystals which decomposed without melting.

15 Example 39

By the application of the above described technique and related procedures known to those skilled in the art and using the appropriate chroman-3-ylamine hydrochlorides, the following compounds were prepared:

20 (*R,S*)-5-(2-aminoethyl)-1-(6-hydroxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound **31**, table 1)

Example 40

25 2-[3-(2,2-Dimethyl[1,3]dioxolan-4-yl)propyl]isoindole-1,3-dione

To a stirred solution of 3-(2,2-dimethyl-[1,3]dioxolan-4-yl)propylamine (1.05 g, 6.60 mmol) and carboethoxyphthalimide (1.45 g, 6.60 mmol) in acetonitrile (10 mL) at room temperature was added triethylamine (0.92 mL, 6.60 mmol) in one portion and the resulting mixture was stirred at room temperature for 18 hours, evaporated in vacuo and the residue was dissolved in ethyl acetate (50 mL). The solution was washed with brine, 10% citric acid solution and brine, then dried over anhydrous magnesium sulfate. Filtration and concentration in vacuo gave an oil which was purified by column

chromatography on silica using ethyl acetate – petroleum ether mixture as eluent to afford a colourless oil.

Example 41

5

2-(4,5-Dihydroxypentyl)isoindole-1,3-dione

To a stirred solution of 2-[3-(2,2-dimethyl[1,3]dioxolan-4-yl)propyl]isoindole-1,3-dione (1.65 g, 5.70 mmol) in THF (20 mL) at room temperature was added 2N HCl solution (15 mL, 30 mmol) in one portion and the resulting mixture was stirred at room temperature for two hours and then evaporated in vacuo to half of the initial volume. The residue was saturated with NaCl and extracted with ethyl acetate. The organic phase was dried by anhydrous magnesium sulfate. Filtration and concentration in vacuo afforded a colourless oil.

15

Example 42

By the application of the technique described in example 8 to 2-(4,5-dihydroxypentyl)isoindole-1,3-dione, the following compound was prepared:

2-[5-(tert-Butyldimethylsilanyloxy)-4-hydroxypentyl]isoindole-1,3-dione

20

Example 43

By the application of the technique described in example 11 to 2-[5-(tert-butyldimethylsilanyloxy)-4-hydroxypentyl]isoindole-1,3-dione, the following compound was prepared:

2-[5-(tert-Butyldimethylsilanyloxy)-4-oxopentyl]isoindole-1,3-dione

Example 44

(S)-5-(3-aminopropyl)-1-(5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride (compound 5, table 1)

30

A stirred mixture of (S)-5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl amine hydrochloride (0.22 g, 1.0 mmol), 2-[5-(tert-butyldimethylsilanyloxy)-4-oxopentyl]isoindole-1,3-dione (0.38 g, 1.05 mmol), potassium thiocyanate (0.11 g, 1.10 mmol), water (0.18 g, 1.0 mmol) and acetic acid (0.3 mL, 5.0 mmol) in ethyl acetate (3

mL) was refluxed for 7 hours, cooled to room temperature, washed by sodium bicarbonate solution, dried over anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.18 g) was dissolved in a mixture of isopropanol (5 mL) and THF (2 mL). Water (0.8 mL) and sodium borohydride (0.066 g, 1.74 mmol) were added at room temperature and the mixture was stirred for 1.5 hours. Acetic acid (0.6 mL, 10 mmol) was added and the solution was refluxed for two hours then evaporated in vacuo to dryness. The residue was taken up into acetone, the solid was filtered off, and the filtrate was acidified with 2N HCl solution in ethyl acetate. The precipitate was collected and washed with acetone to afford crystals, which decomposed without melting.

Example 45

15 (*R*)-5-(3-aminopropyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride

A stirred mixture of (*R*)-6,8-difluorochroman-3-ylamine hydrochloride (0.11 g, 0.50 mmol), 2-[5-(*tert*-Butyldimethylsilyloxy)-4-oxopentyl]isoindole-1,3-dione (0.19 g, 0.55 mmol), potassium thiocyanate (0.055 g, 0.55 mmol), water (0.009 g, 0.50 mmol) and acetic acid (0.15 mL, 2.5 mmol) in ethyl acetate (1.5 mL) was refluxed for 7 hours, cooled to the room temperature, washed by sodium bicarbonate solution, dried over anhydrous magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.10 g) was dissolved in the mixture of isopropanol (2.5 mL) and THF (1 mL). Water (0.4 mL) and sodium borohydride (0.038 g, 1.0 mmol) were added at room temperature and the mixture was stirred for 1.5 hours. Acetic acid (0.3 mL, 5 mmol) was added and the solution was refluxed for two hours and evaporated in vacuo to dryness. The residue was taken up in acetone, the solid was filtered off, and the filtrate was acidified with 2N HCl solution in ethyl acetate. The precipitate was collected and washed with acetone to afford crystals, which decomposed without melting.


Example 46

(*R,S*)-5-(3-aminopropyl)-1-(6-hydroxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride

- 5 A stirred mixture of 6-hydroxythiochroman-3-ylamine hydrochloride (0.22 g, 1.0 mmol), 2-[5-(tert-Butyldimethylsilanyloxy)-4-oxopentyl]isoindole-1,3-dione (0.38 g, 1.05 mmol), potassium thiocyanate (0.11 g, 1.10 mmol), water (0.18 g, 1.0 mmol) and acetic acid (0.3 mL, 5.0 mmol) in ethyl acetate (3 mL) was refluxed for 7 hours, cooled to room temperature, washed by sodium bicarbonate solution, dried over anhydrous
- 10 magnesium sulphate and evaporated in vacuo. The residue was purified by column chromatography on silica using ethyl acetate – petroleum ether mixture as eluent. The resulting oil (0.17 g) was dissolved in the mixture of isopropanol (5 mL) and THF (2 mL). Water (0.8 mL) and sodium borohydride (0.066 g, 1.74 mmol) were added at room temperature and the mixture was stirred for 1.5 hours. Acetic acid (0.6 mL, 10 mmol)
- 15 was added and the solution was refluxed for two hours and evaporated in vacuo to dryness. The residue was taken up into acetone, the solid was filtered off and the filtrate was acidified with 2N HCl solution in ethyl acetate. The precipitate was collected and washed with acetone to afford crystals, which decomposed without melting.

Claims

1. A compound of formula I:



I


- where R_1 , R_2 and R_3 are the same or different and signify hydrogens, halogens, alkyl, alkyloxy, hydroxy, nitro, amino, alkylcarbonylamino, alkylamino or dialkylamino group;
- 10 R_4 signifies hydrogen, alkyl or alkylaryl group; X signifies CH_2 , oxygen atom or sulphur atom; n is 1, 2 or 3, with the proviso that when n is 1, X is not CH_2 ; and the individual (R)- and (S)-enantiomers or mixtures of enantiomers and pharmaceutically acceptable salts thereof; wherein the term alkyl means hydrocarbon chains, straight or branched, containing from one to six carbon atoms, optionally substituted by aryl, alkoxy, halogen,
- 15 alkoxy carbonyl or hydroxycarbonyl groups; the term aryl means a phenyl or naphthyl group, optionally substituted by alkyloxy, halogen or nitro group; the term halogen means fluorine, chlorine, bromine or iodine.
2. A compound according to claim 1, comprising: (S)-5-(2-aminoethyl)-1-(1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (S)-5-(2-aminoethyl)-1-(5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-chroman-3-yl-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-(8-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-(6-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-(8-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (R)-5-(2-aminoethyl)-1-(6-fluorochroman-3-yl)-1,3-dihydroimidazole-2-thione

hydrochloride; (*R*)-5-(2-aminoethyl)-1-(8-fluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6,7-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*S*)-5-(2-aminoethyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6,7,8-trifluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6-chloro-8-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6-methoxy-8-chlorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6-nitrochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(8-nitrochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-[6-(acetylamino)chroman-3-yl]-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-aminomethyl-1-chroman-3-yl-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-aminomethyl-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-aminoethyl)-1-(6-hydroxy-7-benzylchroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-aminomethyl-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(3-aminopropyl)-1-(6,8-difluorochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*S*)-5-(3-aminopropyl)-1-(5,7-difluoro-1,2,3,4-tetrahydronaphthalen-2-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R,S*)-5-(2-aminoethyl)-1-(6-hydroxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R,S*)-5-(2-aminoethyl)-1-(6-methoxythiochroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-benzylaminoethyl)-1-(6-methoxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-5-(2-benzylaminoethyl)-1-(6-hydroxychroman-3-yl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-1-(6-hydroxychroman-3-yl)-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione hydrochloride; (*R*)-1-(6,8-difluorochroman-3-yl)-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione hydrochloride or (*R*)-1-chroman-3-yl-5-(2-methylaminoethyl)-1,3-dihydroimidazole-2-thione hydrochloride.


30

3. A process for the preparation of the individual (*R*)- and (*S*)-enantiomers or mixtures of enantiomers and pharmaceutically acceptable salts of a compound of

formula I of claim 1 which comprises reacting the individual (R)- or (S)-enantiomers or mixtures of enantiomers of a compound of Formula II







5 where X is CH₂, oxygen or sulphur; R₁, R₂ and R₃ are the same or different and signify hydrogens, halogens, alkyl, alkyloxy, hydroxy, nitro, alkylcarbonylamino, alkylamino or dialkylamino group; with a compound of formula III



10

where n signifies 1, 2 or 3; when n is 1 or 2, R₄ signifies hydrogen, alkyl group; R₅ signifies a hydroxyl protecting group and R₆ signifies an amino protecting group; when n signifies 3, R₅ is defined as above but R₄ and R₆ taken together represent a
 15 phthalimido group; and with a water soluble thiocyanate salt in the presence of an organic acid in a substantially inert solvent, followed by subsequent deprotection of the intermediate products IV-VII:



wherein the water soluble thiocyanate salt is an alkali metal thiocyanate salt or a tetraalkylammonium thiocyanate salt, and wherein the term alkyl means hydrocarbon chains, straight or branched, containing from one to six carbon atoms, optionally substituted by aryl, alkoxy, halogen, alkoxycarbonyl or hydroxycarbonyl groups; the term aryl means a phenyl or naphthyl group, optionally substituted by alkyloxy, halogen or nitro group; the term halogen means fluorine, chlorine, bromine or iodine.

10 4. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1 or 2, in combination with a pharmaceutically effective carrier.

5. A compound according to claim 1 or 2, for use as a medicament.

15


6. The use of a compound according to claim 1 or 2, in the manufacture of a medicament for treating disorders where a reduction in the hydroxylation of dopamine to noradrenaline is of therapeutic benefit.

7. The use of a compound according to claim 1 or 2, in the manufacture of a medicament for treating a subject afflicted by cardiovascular disorders.

8. The use of a compound according to claim 1 or 2, in the manufacture of a medicament for treating hypertension or chronic heart failure.

9. The use of a compound according to claim 1 or 2, in the manufacture of a medicament for use in inhibiting dopamine- β -hydroxylase.


10. A compound of formula III



15 where n signifies 2, R_4 signifies hydrogen or alkyl group; R_5 signifies a hydroxyl protecting group and R_6 signifies an amino protecting group, wherein the hydroxyl protecting group is a trialkylsilyl, triphenylsilyl, phenyldialkylsilyl or alkyl diphenylsilyl group; and the amino protecting groups is an alkyl carbamate or alkylaryl carbamate group, and wherein the term alkyl means hydrocarbon chains, straight or branched,
 20 containing from one to six carbon atoms, optionally substituted by aryl, alkoxy, halogen, alkoxy carbonyl or hydroxy carbonyl groups; the term aryl means a phenyl or naphthyl group, optionally substituted by alkoxy, halogen or nitro group; the term halogen means fluorine, chlorine, bromine or iodine.

25


11. A compound of formula III



where n signifies 3, R₅ signifies a hydroxyl protecting group; and R₄ and R₆ taken together represent a phthalimido group, wherein the hydroxyl protecting group is a trialkylsilyl, triphenylsilyl, phenyldialkylsilyl or alkyldiphenylsilyl group.

5

12. A compound of formula II



II

wherein R₁, R₂ and R₃ may be the same or different and signify fluorine or hydrogen
10 with the proviso that at least one R₁, R₂ and R₃ signifies fluorine where X is the same as defined in claim 1.

13. (R)-6,8-difluoro-3,4-dihydro-2H-1-benzopyran-3-amine.

15 14. A compound substantially herein described with reference to and as shown in the accompanying examples.

15. A pharmaceutical composition substantially herein described with reference to and as shown in the accompanying examples.


20

16. A process substantially herein described with reference to and as shown in the accompanying examples.

25

ABSTRACT**PERIPHERALLY-SELECTIVE INHIBITORS OF DOPAMINE- β -HYDROXYLASE**
AND METHOD OF THEIR PREPARATION

5 Compounds of formula I and a method for their preparation are described:



I

where X is CH₂, O or S, and n is 1, 2 or 3, with the proviso that if X is CH₂, n is not 1.

10

The compounds have potentially valuable pharmaceutical properties for the treatment of cardiovascular disorders such as hypertension and chronic heart failure.

